



South Australian
Certificate of Education

Specialist Mathematics

2023

Question booklet 1

Questions 1 to 7 (55 marks)

- Answer **all** questions
- Write your answers in this question booklet
- You may write on page 16 if you need more space
- Allow approximately 65 minutes
- Approved calculators may be used — complete the box below

Examination information

Materials

- Question booklet 1
- Question booklet 2
- Formula sheet
- SACE registration number label

Instructions

- Show appropriate working and steps of logic in the question booklets
- State all answers correct to three significant figures, unless otherwise instructed
- Use black or blue pen
- You may use a sharp dark pencil for diagrams

Total time: 130 minutes

Total marks: 100

© SACE Board of South Australia 2023

Attach your SACE registration number label here

Graphics calculator

1. Brand _____

Model _____

2. Brand _____

Model _____



Government
of South Australia

SACE
BOARD
OF SOUTH
AUSTRALIA

Question 1 (5 marks)

Consider the plane $P: x + 2y - z = 4$ and the line l defined by the following parametric equations, where t is a real parameter.

$$\begin{cases} x = 1 + t \\ y = 2 - 2t \\ z = 3 - t \end{cases}$$

(a) Show that the point of intersection of P and l is $A(0, 4, 4)$.

(3 marks)

(b) Find the equation of the plane which passes through A and is perpendicular to l .

(2 marks)

(ii) Hence show that $\frac{d\theta}{dt} = \left(\frac{50}{2500 + x^2} \right) \frac{dx}{dt}$.

(2 marks)

(c) Hence find the rate of change, $\frac{d\theta}{dt}$, when $t = 9$ seconds. Give the answer correct to three significant figures.

(3 marks)

Question 4 (7 marks)

(a) Prove by mathematical induction that $4^n + 15n - 1$ is divisible by 9 for all positive integers n .



(5 marks)

Question 6 (10 marks)

The population B of butterflies in a butterfly house has a rate of growth, $\frac{dB}{dt}$, that is modelled by the differential equation

$$\frac{dB}{dt} = 0.1B \left(1 - \frac{B}{K} \right),$$

where K is a positive constant and t is the time in months, $t \geq 0$, and B equals the number of butterflies in the population.



Source: © Jen567 | Pixabay.com

(a) (i) Show that $\frac{K}{B(K-B)} = \frac{1}{B} + \frac{1}{K-B}$.

(1 mark)

(ii) Using integration techniques, show that the butterfly population can be modelled by


$$B = \frac{K}{1 + Ae^{-0.1t}}$$

for some constant A .

(4 marks)

Question 7 (10 marks)


- (a) Given that $\int xe^x dx = xe^x - e^x + c$, use integration by parts to show that $\int x^2 e^x dx = e^x(x^2 - 2x + 2) + k$, where c and k are constants.



(2 marks)

- (b) Consider the functions $f(x) = \sqrt{x}$ and $g(x) = \frac{e^x}{4}(4 - x^2)$.

- (i) Show that $f(g(x)) = \frac{1}{2}\sqrt{e^x(4 - x^2)}$.



(1 mark)

- (ii) On Figure 6 below draw the graph of $y = f(g(x)) = \frac{1}{2}\sqrt{e^x(4 - x^2)}$ for $-2 \leq x \leq 2$.





Figure 6

(3 marks)

(c) The curve $y = f(g(x))$ sketched in Figure 6 is rotated 2π radians about the x axis for $-2 \leq x \leq 2$ to form a solid.

(i) Show that the volume of the solid formed by the rotation is given by

$$V = \frac{\pi}{4} \int_{-2}^2 e^x (4 - x^2) dx.$$



(1 mark)

(ii) Using part (a), show that the volume of the solid is $\frac{\pi}{2}(e^2 + 3e^{-2})$.



(3 marks)

You may write on this page if you need more space to finish your answers to any of the questions in this question booklet. Make sure to label each answer carefully (e.g. 5(a)(ii) continued).

A large grid of graph paper, consisting of 20 columns and 30 rows of small squares, intended for writing answers to questions.



South Australian
Certificate of Education

Specialist Mathematics

2023

Question booklet 2

Questions 8 to 10 (45 marks)

- Answer **all** questions
- Write your answers in this question booklet
- You may write on page 11 if you need more space
- Allow approximately 65 minutes
- Approved calculators may be used — complete the box below

2

© SACE Board of South Australia 2023

Copy the information from your SACE label here

SEQ	FIGURES	CHECK LETTER	BIN
<input type="text"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	<input type="text"/>	<input type="text"/>

Graphics calculator

1. Brand _____

Model _____

2. Brand _____


Model _____



Government
of South Australia


(c) It may be assumed that a normal vector to P_1 and P_2 is $l = [2, 2, -1]$.

(i) Find $|\vec{BA} \times l|$.




(2 marks)

(ii) Find $|\vec{AT}|$, using the result of part (a).




(2 marks)

(iii) Find the equation of the line normal to P_2 through $B(-1, -6, -2)$.



(2 marks)

(iv) Show that the line found in part (c)(iii) intersects P_1 at $T(3, -2, -4)$.



(2 marks)

Question 9 (15 marks)

On the Argand diagram in Figure 10 is the set of complex numbers z such that $z = \text{cis}\theta$ for $-\pi < \theta \leq \pi$.
 The Argand diagram in Figure 11 shows the complex numbers z_1, z_2, z_3, z_4, z_5 which are the zeros of the polynomial $z^5 - 1$.




Figure 10




Figure 11

(a) Using De Moivre's theorem or otherwise, write the zeros of $z^5 - 1$ in exact polar form.

(3 marks)

(b) Consider the function $f(z) = z^2 + z + 1 + \frac{1}{z} + \frac{1}{z^2}$, where $z = \text{cis}\theta$.

(i) Show that $|f(z)| \leq 5$.


(2 marks)

(ii) Find a value of z for which $|f(z)| = 5$.

(1 mark)

Question 10 (15 marks)

(a) Show that $\int \cos^2 x \, dx = \frac{1}{2}x + \frac{1}{4}\sin 2x + c$, where c is a constant.



(2 marks)

(b) On Figure 13, draw the curve defined by the following parametric equations, where t is time in seconds.

$$\begin{cases} x = 4\sqrt{2} \sin t \\ y = \frac{\sqrt{2}}{2} \sin 2t \end{cases} \text{ for } 0 \leq t \leq 2\pi$$





Figure 13


(3 marks)

(iii) Hence show that the exact distance travelled by the dragonfly over the interval $\frac{\pi}{4} \leq t \leq \frac{3\pi}{4}$ is $\sqrt{2}(2\pi - 1)$.



(3 marks)

You may write on this page if you need more space to finish your answers to any of the questions in this question booklet. Make sure to label each answer carefully (e.g. 10(c)(ii) continued).



SPECIALIST MATHEMATICS FORMULA SHEET

Circular functions

$$\sin^2 A + \cos^2 A = 1$$

$$\tan^2 A + 1 = \sec^2 A$$

$$1 + \cot^2 A = \operatorname{cosec}^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$= 2 \cos^2 A - 1$$

$$= 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$2 \sin A \cos B = \sin(A + B) + \sin(A - B)$$

$$2 \cos A \cos B = \cos(A + B) + \cos(A - B)$$

$$2 \sin A \sin B = \cos(A - B) - \cos(A + B)$$

$$\sin A \pm \sin B = 2 \sin \frac{1}{2}(A \pm B) \cos \frac{1}{2}(A \mp B)$$

$$\cos A + \cos B = 2 \cos \frac{1}{2}(A + B) \cos \frac{1}{2}(A - B)$$

$$\cos A - \cos B = -2 \sin \frac{1}{2}(A + B) \sin \frac{1}{2}(A - B)$$

Matrices and determinants

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then $\det A = |A| = ad - bc$ and


$$A^{-1} = \frac{1}{|A|} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Measurement

Area of sector, $A = \frac{1}{2} r^2 \theta$, where θ is in radians.

Arc length, $l = r\theta$, where θ is in radians.

In any triangle ABC :



$$\text{Area of triangle} = \frac{1}{2} ab \sin C$$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Quadratic equations

$$\text{If } ax^2 + bx + c = 0 \text{ then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Distance from a point to a plane

The distance from (x_1, y_1, z_1) to

$Ax + By + Cz + D = 0$ is given by

$$\frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}.$$

Derivatives

$f(x) = y$	$f'(x) = \frac{dy}{dx}$
$\arcsin x$	$\frac{1}{\sqrt{1-x^2}}$
$\arccos x$	$\frac{-1}{\sqrt{1-x^2}}$
$\arctan x$	$\frac{1}{1+x^2}$

Properties of derivatives

$$\frac{d}{dx} (f(x)g(x)) = f'(x)g(x) + f(x)g'(x)$$

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

$$\frac{d}{dx} f(g(x)) = f'(g(x))g'(x)$$

Arc length along a parametric curve

$$l = \int_a^b \sqrt{\mathbf{v} \cdot \mathbf{v}} dt, \text{ where } a \leq t \leq b.$$

Integration by parts

$$\int f'(x)g(x) dx = f(x)g(x) - \int f(x)g'(x) dx$$

Volumes of revolution

About x axis, $V = \int_a^b \pi y^2 dx$, where y is a function of x .

About y axis, $V = \int_c^d \pi x^2 dy$, where y is a one-to-one function of x .